#### **General Disclaimer**

### One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some
  of the material. However, it is the best reproduction available from the original
  submission.

Produced by the NASA Center for Aerospace Information (CASI)

In the under NASA and the contract and wide disfic. An Resources Survey pion and without lish and the contract of the contra

"Made available under NASA sponsorship in the interest of early and wide discomination of Earth Resources Survey communication and without liability and use made thereof?" 亚

FIFTH QUARTERLY (TYPE II) PROGRESS REPORT 27.7-10.0.45.

CR-149181

Project Title: Development of a Multi-disciplinary

ERTS User Program in the State of Ohio

Ref. 20900

Contract Number: NAS5-22399

Principal Investigator: Paul E. Baldridge

Deputy Director for Community Services
Department of Economic and Community Development

P.O. 1001

Columbus, Ohio 43216

September 10, 1976

(E77-10045) DEVELOPMENT OF A

MULTI-DISCIPLINARY ERTS USER PROGRAM IN THE

STATE OF OHIO Quarterly Progress Report
(Ohio Dept. of Economic and Community) 5 p

HC A02/MF A01

CSCL 05B G3/43 00045

Prepared for

GODDARD SPACE FLIGHT CENTER Greenbelt, Maryland 20771

20900

Scientific Investigation Support, Code 902.6 Contract No. NAS5-22399 ERTS Investigation No. 20900 Items No. 1 and 2

RECEIVED

NOV 18 1976

SIS/902.6

## Section A - Problems

- 1. Statewide Land Use Inventory no problems
- 2. Socio-economic merging with LANDSAT data. This section of work is still waiting on results of the Statewide Land Use Inventory.
- 3. Services Development. The problems at this time are associated with the Economic Development Forestry Resource Management portion where small-scale aircraft coverage over the test site has not yet been satisfactorily accomplished; also the Lake Erie Sedimentation study has some difficulties with the correlation of sedimentation ground observation data and the LANDSAT sedimentation patterns.

## Section B - Accomplishments

- 1. Statewide Land Use Inventory All of the geographic boundaries have been digitized at this point. In addition, the computer program to convert Bendix tape output format to OCAP format has been complete; and debugged. Bendix has categorized the state as a whole and the only remaining task is to geometrically correct and map into the U.S.G.S. seven and one-half minute quadrangles.
- Socio-economic merging with Landsat data. Accomplishment all stated in previous reports.
- 3. Service Development -

Urban Land Use Analysis:

Levels I and II land use information has been delineated from six frames of IR color transparencies which were taken by NASA over Columbus on August 12, 1975. Land use class information was transferred to black and white acetate topographic base maps at a scale of 1:24,000. The entire county is actually covered by 18 base maps, some showing only small areas at the edges or corners of the county with few land use categories. The major metropolitan area of Columbus, however, is covered by four quad sheets, covering

some 231 square miles (590 km)<sup>2</sup> of the 407-square mile county area. Following the USGS land use classification system (with some modification), the following categories have been delineated:

- 1.1 Residential (single and multiple dwellings)
- 1.2 Commercial and services (shopping centers, warehouses, etc.)
- 1.3 Industrial (manufacturing plants)
- 1.4 Extractive (quarries)
- 1.5 Transportation (roads, railroads, airports)
- 1.6 Institutional (schools, churches, research institutes, hospitals, etc.)
- 1.7 Recreation, public and private (parks, golf courses, etc.)
- 1.8 Mixed
- 1.9 New construction

Each land use category has been color coded. Some difficulty has been experienced in the identification of industrial and commercial sites.

Larger scale (1:40,000) imagery flown by NASA over Columbus in 1973 was used to differentiate between small industrial and commercial activities which could not be properly interpreted from the high-altitude data.

\* All mineral surface mining and exploration:

Several overlays showing linear features derived from LANDSAT data have been completed. Several techniques were used to detect, trace, and code linears. Most analysis was done using a 200 line per inch Ronchi grating in conjunction with a projection system. More than 300 linears varying in length from 1 to 20 miles have been plotted on 1:250,000-scale maps of Ohio.

### Section C - Significant Results

- 1. Statewide Land Use Inventory None to report at this time.
- 2. Socio-economic merge none to report at this time.
- 3. Service Development Urban Land Use Analysis -

A preliminary comparison of the land use maps derived from the highaltitude photography and imagery generated from land use CCT's at Battelle shows considerable correlation of both data. The degree of success of interpreting LANDSAT data, however, is directly related to the P.I.'s ability to correlate selected area samples recorded on highaltitude data. Some land use classes, however, such as areas covered by institutional and commercial features always require identification from data with a better resolution than obtainable with LANDSAT data. Most land use features are best identified on LANDSAT data taken over a full year, showing the effects of the four seasons. It thus appears that the most cost effective and timely way of providing land use data to end users is to use LANDSAT data for identifying 75 to 90 percent of all land use categories, using temporal data. The remainder of the urban area land users may then be interpreted using primarily high flight data and, in selected cases, conventional aircraft or reference data.

All mineral surface mining and exploration -

Close examination of LANDSAT, Skylab and aircraft data covering the same area where linears were found seems to indicate that the type of resolution characteristic of LANDSAT data results in a better definition of linears of l or more miles. Long-line linears have been found in the vicinity of known gas production areas in Ohio.

# Section D - Publications

 The Ohio Land Allocation Model: Report on Phase II by Oscar Fisch and Steven I. Gordon submitted to the Department of Economic and Community Development July, 1976.

# Section E - Recommendations

1. None.